# Antimicrobial Activity of Lactic Acid Bacteria Found in Dadiah on Disease-Causing Skin Infections

by Dessy Abdullah

Submission date: 27-May-2022 04:35PM (UTC+0800)

**Submission ID:** 1845220872

File name: ound\_in\_Dadiah\_on\_Disease-Causing\_Skin\_Infections\_compressed.pdf (441.32K)

Word count: 3865
Character count: 24405



Volume 15

Number 2

April-June 2021



# Indian Journal of Forensic Medicine & Toxicology

Website: www.ijfmt.com

### Indian Journal of Forensic Medicine & Toxicology

### **EDITOR** in Chief

### Prof. R K Sharma

Formerly at All India Institute of Medical Sciences, New Delhi, E-mail: editor.ijfmt@gmail.com

#### **EDITOR**

#### Prof. Dr. Adarsh Kumar

Forensic Medicine & Toxicology, AIIMS, New Delhi

### INTERNATIONAL EDITORIAL ADVISORY BOARD

- 1. Prof Mete Gulmen Cukurova University, TURKEY
- Prof. Leandro Duarte De Carvalho, Minas Gerais, Belo Horizante, Brazil
- Prof. Donata Favretto (Full Professor) Forensic Toxicology at University of Padova. Italy
- Prof. Babak Mostafazadeh Department of Forensic Medicine & Toxicology, Shahid Beheshti University of Medical Sciences, Tehran-Iran
- 5. Prof Halis Dokgoz, Mersin University, TURKEY
- 6. Prof Jozef Sidlo, Comenius University, Bratislava, SLOVAKIA
- Dr. Rahul Pathak (Lecturer) Forensic Science, Dept of Life Sciences Anglia Ruskin University, Cambridge, United Kingdom
- Dr. Hareesh (Professor & Head) Forensic Medicine, Ayder Referral Hospital, College of Health Sciences, Mekelle University, Mekelle Ethiopia East Africa
- Dr. Mokhtar Ahmed Alhrani (Specialist) Forensic Medicine & Clinical Toxicology, Director of Forensic Medicine Unit, Attorney General's Office, Sana'a, Yemen
- Dr. Sarathchandra Kodikara (Senior Lecturer) Forensic Medicine, Department of Forensic Medicine, Faculty of Medicine, University of Peradeniya, Sri Lanka
- Dr Noha A. Magdie El Rafie, Forensic Toxicology, Ain Shams University, Cairo, EGYPT

### **SCIENTIFIC COMMITTEE**

- Prof Udai Pratap Singh, Department of Anthropology Lucknow University Lucknow
- Dr Anil Rahule (Associate Professor) Dept of Anatomy, Govt Medical College Nagpur
- Dr Shankar Bakkanwar (Associate Professor) Forensic Medicine, Kasturba Medical College, Manipal, Karnatakad
- 4. Dr K. Ravikumar Raksha Shakti University, Ahmedabad, Gujrat.
- Dr. Pragnesh Parmar (Associate Professor) Forensic Medicine, Valsad, Gujrat
- Dr Vandana Mudda (Awati) (Associate Prof) Dept of FMT, M.R.Medical College,Gulbarga, Karnataka,
- Dr. Asha Srivastava (Senior Scientific Officer) Forensic Psychology, Central Forensic Science Laboratory, CBI, Delhi
- Dr. Lav Kesharwani (Asst.Prof.) School of Forensic Science, Sam Higginbottom Institute of Agriculture Technology & Sciences, Allahabad U.P.
- Dr. Anu Sharma (Associate Prof) Dept of Anatomy, DMCH, Ludhiana (PB)
- Dr. Shalini Gupta ( Prof ) Oral Pathology and Microbiology, Dental Sciences King George Medical University, Lucknow, UP
- 11. Dr Rituja Sharma, Associate Prof, Law Banasthali Vidyapeeth Jaipur

"Indian Journal of Forensic Medicine & Toxicology" is peer reviewed quarterly journal. It deals with Forensic Medicine, Forensic Science, Toxicology, DNA fingerprinting, sexual medicine and environment medicine. It has been assigned International standard serial No. p-9973-9122 and e-0973-9130. The Journal has been assigned RNI No. DELENG/2008/21789. The journal is indexed with Index Copemicus (Poland) and is covered by EMBASE (Excerpta Medica Database). The journal is also abstracted in Chemical Abstracts (CAS) database (USA. The journal is also covered by EBSCO (USA) database. The Journal is now part of UGC, DST and CSIR Consortia. It is now offical publication of Indian Association of Medico-Legal Experts (Regd.).

### NATIONAL EDITORIAL ADVISORY BOARD

Prof Sudhir K Gupta - Head, Department of Forensic Medicine All India Institute of Medical Sciences, New Delhi

#### Members

- 1. Prof. SK Dhattarwal, Forensic Medicine, PGIMS, Rohtak, Haryana
- 2. Prof. N K Aggrawal Forensic Medicine, UCMS, Delhi
- Prof Ajay Ghangale Forensic Medicine Dr DY Patil Medical College, Pune Meharachtra
- Dr. Amar Jyoti Patwory Professor, Forensic Medicine NEIGRIHMS, Shillong
- Dr S. Venkata Raghava Professor, Forensic Medicine, Banglore Medical College, Bengaluru
- Prof Praveen Arora, Professor Department of Forensic Medicine & Toxicology, SAIMS, Indore
- Dr. Pankaj Datta (Principal & Head) Department of Prosthodontics, Indraprastha Dental College & Hospital, Ghaziabad
- Dr. Mahindra Nagar (Head) Department of Anatomy, UCMS & GTB Hospital, Delhi
- Dr. Virender Kumar Chhoker Professor Forensic Medicine and Toxicology, Santosh Medical College, Ghaziabad, UP
- Dr. Dayanand G Gannur (Professor) Department of Forensic Medicine & Toxicology, Shri BM Patil Medical College, Hospital & Research centre, Bijapur, Karnataka
- Dr. Alok Kumar Professor Department of Forensic Medicine & Toxicology, UP Rural Institute of Medical Sciences and Research, Saifai, Etawah, U.P.

### Print-ISSN:0973-9122 Electronic - ISSn: 0973-9130

Frequency. Quarterly, © All Rights reserved The views and opinions expressed are of the authors and not of the Indian Journal of Forensic Medicine & Toxicology, Indian Journal of Forensic Medicine & Toxicology does not guarantee directly or indirectly the quality or efficacy of any products or service featured in the advertisement in the journal, which are purely commercial.

### Website: www.ijfmt.com

### Editor

### Dr. R.K. Sharma

Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall, Sector- 32, Noida
- 201 301 (Uttar Pradesh)

### Printed, published and owned by

### Dr. R.K. Sharma

Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall, Sector- 32, Noida

- 201 301 (Uttar Pradesh)

### Published at

### Institute of Medico-legal Publications

Logix Office Tower, Unit No. 1704, Logix City Centre Mall, Sector- 32, Noida - 201 301 (Uttar Pradesh)



### Indian Journal of Forensic Medicine & Toxicology

Contents

Volume 15. Number 2

April-June 2021

Volume 15, Number 2	
1.	Psychological Aspects in Terms of Physical, Mental and Sociological Factors During Worldwide  Corona Crisis
2.	Anatomical Variation of Superficial Branch of Radial Nerve in Gujrat Region
3.	Combined Effect of Neural Tissue Mobilization and Deep Friction Massage in Piriformis  Syndrome: A Research Protocol
4.	Police Brutality and Custodial Torture in Technological Era: Need for Anti-Torture Law in India - A Critical Analysis
5.	Study of Socio-Demographic Profile of Thoraco-Abdominal Injuries in Fatal Road Traffic Accidents in Mangalore City
6.	Study of Mortality Due to Thoraco-Abdominal Injuries in Road Traffic Accidents in Mangalore City 32 Abymon K K, Mohd Amjad Bhat, Azia Manzoor Bhat
7.	Pattern of Thoraco-Abdominal Injuries in Fatal Road Traffic Accidents in Mangalore City
8.	An Ayurvedic Approach for Treatment of Fissure in Ano: A Case Report of 32 Year Old Patient43 Alok kumar Diwedi, Kiran Khandare, Rakesh Khatana, Harjodh Singh
9.	Ozone Therapy - A Boon to Dental Sciences
10.	Hyperammonaemia in Absence of Deranged Liver Enzymes with Valproate: A Case Report
11.	Assessment of Anxiety during COVID-19 Pandemic among General Population of India
12.	A Conceptual Study on Prevention & Management of Covid -19 Pandemic with Special Reference to  Janapadodwamsa

487.	The Use of Pentoxifylline Topical in the Treatment of Alopecia in Rats	3241
488.	Study of Anti-Ovarian Antibody, Anti-FSH and Anti-LH Antibodies Along with Their Receptors in Polycystic Ovarian Syndrome	3250
489.	Correlation between Salivary Cotinine Levels and Cigarette Smoking with Recurrent Aphthous Stomatitis	3258
490.	Contaminant Investigations of Regularly used Eye Contact Lenses	3264
491.	Viral Profile and Clinical Characteristic in Acute Asthma Exacerbation Patients	3276
492.	Capillary Lactate Level in Non-Severe and Severe Community-Acquired Pneumonia Patients  Resti Yudhawati, Kowiy Akbar	3284
493.	Reconstruction of Prevention and Handling of Stunting Policy in Public Health Center	3292
494.	Determinants of Loss to Follow-Up Among MDR-TB Patients	3297
495.	Motivational Interviewing and Physical Activity on Quality of Life of Type 2 DM Patients in Makassar City  **Ridwan Amiruddin, Nurhaedar Jafar, Jumriani Ansar, Uliadi Barrung Limbong, Risnah**	3303
496.	Antimicrobial Activity of Lactic Acid Bacteria Found in Dadiah on Disease-Causing Skin Infections	3314
497.	Children and Stunting Phenomenon in Indonesia: Legal Perspective	
498.	Which Predisposing Factors Influence The Farmers in Compliance of Personal Protective Equipment (PPE)?	3323
499.	Method Development and Optimization for Colchicine Estimation in Toxicological Analysis using Dispersive Liquid-Liquid Micro-Extraction and HPLC-UV	3328

## Antimicrobial Activity of Lactic Acid Bacteria Found in Dadiah on Disease-Causing Skin Infections

### Rinita Amelia<sup>1</sup>, Dessy Abdullah<sup>2</sup>, Yudha E. Pratama<sup>3</sup>, Endang Purwati<sup>4</sup>

<sup>1,2</sup>Doctoral Program Student Lincoln of University College, Faculty of Health and Science, Bharu City Malaysia,
<sup>1,2</sup>Lecture Baiturrahmah University, Medical Faculty, Padang Indonesia, <sup>3</sup>Doctoral Program Student Andalas University, Faculty of Animal Science, Padang, Indonesia, <sup>4</sup>Professor Andalas University, Faculty of Animal Science, Biotechnology/Technology of Product Husbandry, Padang Indonesia

### Abstract

Dadiah is a fermented food made from buffalo milk containing lactic acid bacteria (LAB). The potential of LAB as an antimicrobial organism is expected to inhibit or kill pathogens, especially those causing inflammations in the skin. Dadiah used in this study was obtained from the Lintau area of Tanah Datar District, West Sumatra, Indonesia. LAB isolation and identification involved the use of MRS Broth (MERCR) and MRS Agar (MERCK) media and the pathogenic bacteria used include Staphylococcus aureus, Propionibacterium acnes, Acinetobacter Baumannii, Klebsiella pneumonia, and Pseudomonas aeruginosa. The results of this study obtained a total LAB of 7x10<sup>11</sup> CFU / g and showed the highest antimicrobial activity in the Staphylococcus aureus with a14.53 mm clear zone and 13.18 mm bacteria in the Pseudomonas aeruginosa test bacteria.

Keywords: Antimicrobial dadiah, lactic acid bacteria, skin diseases

### Introduction

Skin diseases are the most common types of diseases in developing countries, especially in those with tropical climate. Therefore, humid and hot air throughout the year is highly suitable for the development of infectious, disease-causing pathogens in the skin. The prevalence of such infections in developing countries ranges between 20-80%. As a developing country, Indonesia and its inhabitants are prone to skin diseases of which its main causes include poor geographical and hygiene conditions. Such conditions enhance the rapid development of pathogens and bacteria. Therefore, the attention and recommendations of appropriate health organizations in the preventive measures against skin infections is presently a necessity. The problem of antibiotic resistance has become a worldwide problem, especially during prevention and cure, that necessitates

Corresponding Author:
Prof. drh. Hj. Endang Purwati MS, PhD
purwati17@yahoo.co.id / purwati17@ansci.unand

viable solutions in form of alternative therapy or treatments<sup>[1]</sup>. These forms of alternative therapy include synthetic antibiotics.

Probiotics, vitamins, and phenolics are a few examples of nutraceuticals. These substances are important for skin protection and health, and they are known to assist in medical treatment of dermatological problems<sup>[2]</sup>. This study aims to discover the role of probiotics (Lactobacillus) as a prevention and treatment for skin infections. Many studies have shown that lactic acid bacteria (LAB) helps to activate commensal bacteria due to its probiotic properties, and gets rid of pathogenic bacteria through the antimicrobial properties derived from either bacteriocin found in LAB itself, or the role of its peptides. Some studies also mention that the role of probiotics in nutrition involves boosting the immunity against skin infections caused by various pathogenic bacteria such as Aeromonas bestiarum, Ichthyophthirius multifiliis, Staphylococcus aureus, Corynebacterium acnes, Streptococcus, etc[3].

Dadiah is a traditional food from West Sumatra made of buffalo milk that undergoes a natural fermentation process in a bamboo tube, Dadiah originally made from Lintau in Tanah Datar, contains probiotics such as L. Plantarum which has proven to be effective in maintaining the balance of intestinal microflora, and contains strong antimicrobial properties<sup>[4]</sup>.

Atopic dermatitis is an inflammatory skin disease occurring often in children and the elderly [5]. The dominant bacterial species found in this skin disease is Staphylococcus aureus. The colonies of this bacteria, when present in the skin, can result in a wide range of infections therein. Therefore, antimicrobial treatment is necessary to prevent such bacterial growth [6]. Many broad-spectrum antibiotics can be used as an inhibitor of Staphylococcus aureus, such as penicillin, doxycycline, tetracycline, clindamycin, etc. However, improper use of an antibiotic results in drug resistance. It is also associated with an increased risk of infection and emergence of bacterial colonies, known as "Multidrug-Resistant Organisms (MDRO)" [7]. These Methicillin resistant Staphylococcus aureus (MRSA) and MDRO are the main causes of nosocomial pathogenic infection worldwide, and require serious treatment, as they increase comorbidity and death due to the infection[8]. A major challenge posed by S. aureus is antimicrobial resistance. After the introduction of the antibiotics penicillin and methicillin into clinical practice, strains of antibiotic-resistant S. aureus (MRSA) were subsequently identified [9].

Therefore, this study was aimed to determine the role of Dadiah as a probiotic that can be used to inhibit bacteria-induced skin diseases, especially the inflammatory infections, as a supporting or future treatment alternative against skin diseases.

### Materials and Methods

### Isolation of Lactic Acid Bacteria

One gram of *Dadiah* sample was placed within test tubes containing 9 ml of MRS broth and diluted up to ten logarithmic (10-10) fold. Furthermore, it was added to test tubes containing 900 ul of MRS broth variations of dilution ratio 10-1, 10-2 and 10-7. Furthermore, 100 ul of the serial dilution of 10-9 was inoculated into the MRS Agar media using the spread method, placed in an anaerobic jar and incubated for 48 hours with a temperature 37°C. After incubation, a single colony of LAB was obtained and hence isolated [10].

### Identification of Lactic Acid Bacteria

Identification of LAB from Dadiah samples was done by placing 1g of the sample in a test tube containing 9 ml MRS Broth diluted up to ten (10-1) logarithmic fold. It was further incubated in anaerobic conditions for 24 hours, after which a 100 uL 10-1 dilution was added to the tube containing 900 uL 1 MRS broth dilution 10-2, and a subsequent dilution of 10-8. Consecutively, 100 uL of the serial dilution of 10-8 was inoculated on the media RS Agar with the method of spread, placed within an anaerobic jar, and incubated for 48 hours at 37 °C. A single (choice random) colony that is round, slippery, and either white or yellowish is characteristic of LAB. Finally, this bacterial colony was transferred via streak method and incubated [10].

### **Antimicrobial Activity**

The antimicrobial resistance testing was carried out using five bacteria samples that cause skin disease. LAB culture (1 mL) was centrifuged at 10,000 rpm for 5 minutes at 27° C, and supernatant volume of 50 ul was used to test for antimicrobial resistance. 0.2% of test bacteria (Staphylococcus aureus, Propionibacterium acnes, Acinetobacter Baumannii, Klebsiella pneumonia, and Pseudomonas aeruginosa) was placed on Nutrient Agar (NA) as a growth media for test bacteria, homogenized, injected by micropipette, and incubated at 37 °C. Clear and circular inhibition zones are measured after 12 hours [11].

### Results and Discussion

### Isolation Lactic Acid Bacteria

Dadiah originating from the Lintau area of Tanah Datar District, West Sumatra, Indonesia is known to contain Lactic Acid Bacteria (LAB). It was isolated with MRS as a specific media for LAB growth and incubated for 48 hours. In this study, the amount of LAB used was 7x1011 CFU / g. This result is higher compared to those obtained, while the total LAB obtained from Payakumbuh Dadiah was 8x108 CFU/g [12], in a research at Solok district, the total LAB obtained was as much as 8x108 CFU/g [13]. This is because Dadiah kicks off its fermentation spontaneously; without the need for a fermentation starter. LAB was also found in buffalo milk obtained from several areas in West Sumatra totaling  $3x10^6$  -  $2x10^8$  CFU/g <sup>[14]</sup>. This is consistent with the standards which state that LAB in probiotic food must be present within the range of  $10^6$  -  $10^8$  CFU / Gram <sup>[15]</sup>.

### Identification of Lactic Acid Bacteria

Incubation was done using MRS Broth and then cultured on MRS media. For 48 hours in 37<sup>0</sup> C incubators, a single colony was obtained through macroscopic observations with round, creat white and convex shapes. Microscopic observations showed

that the bacteria were Gram-positive and rod-shaped. Lactic acid bacteria (LAB), belongs to a group of gram-positive, catalase-negative bacteria, which synthesize lactic acid by the fermentation of carbohydrates. The cells are known as coccus or bacillus, usually arranged in pairs or chains. LAB are facultative anaerobes, non-motile and mesophyllic [16].

### **Antimicrobial Activity**

Antimicrobial testing using Staphylococcus aureus, Propionibacterium acnes, Acinetobacter Baumanni, Klebsiella pneumoniae and Pseudomonas aeruginosa commonly found in skin infections can be seen in Table 1.

Table 1. Diameter Clear Zone

In all to	Bacteria test (mm)				
Isolate	Staphylococcus aureus ATCC	Propionibacterium acnes	Acinetobacter baumannii	Klebsiella pneumoniae	Pseudomonas aeruginosa
3A	14,53	7,6	10,24	10,15	13,18

Testing antimicrobial activity involved a well-known method of random selection of 3A isolates from curd which have the potential to function as probiotics, because they can inhibit the growth of pathogenic bacteria which cause skin diseases. The clear zone produced by lactic acid in table 1 shows the existence of antimicrobial activity from LAB isolated from Dadiah samples obtained from Lintau, Tanah Datar District. The ability of LAB to inhibit pathogenic bacteria is due to the presence of its metabolites such as lactic acid, peroxide, and bacteriocin.

Clear zones with the highest numbers were produced in the *Staphylococcus aureus* ATCC test bacteria (14.53 mm) from *Dadiah* isolates. This clear zone is smaller when compared to the study of which used LAB isolated from tempoyak which inhibits the bacterium *Staphylococcus aureus* which a diameter of 19.3 mm [17].

In *Propionibacterium acnes*, test bacteria obtained a clear zone of 7.6mm, the smallest diameter compared to other test bacteria. This result is smaller when compared

with Lactobacillus casei strain of Shirota bacteria of 11.2 mm and therefore higher in inhibitory power than the L. acidophilus HM1 bacteria of 3.3 mm [18]. The clear zone of Acinetobacter baumannii (10.24 mm) obtained in this study is greater than that of which used the Lactobacillus acidophilus LMEM8 against Acinetobacter baumannii and obtained a diameter of 5.13 mm [19]. The results of this study are more productive than that of in testing the antimicrobial activity of Lactobacillus fermentum MTCC 9748 against Klebsiella pneumoniae NTCC 703603 which resulted in a 9.17 mm clear zone [20]. Moreover, the Pseudomonas aeruhginosa test bacteria (13.18 mm clear zone) was not significantly different compared to the research testing of Lactobacillus spp against Pseudomonas aeruginosa bacteria with a range of clear zones 11-21 mm [21].

### Conclusion

Lactic acid bacteria isolated from *Dadiah* of Lintau origin is probiotic in nature, and it shows a highly significant inhibitory effect against *Staphylococcus* 

aureus (of which the largest clear zone of 14.53mm was obtained), Propionibacterium acnes, Acinetobacter baumannii, Klebsiella pneumoniae and Pseudomonas aeruginosa. However, further research is necessary in order to ascertain other probiotic, antimicrobial candidates which inhibit pathogenic bacteria which cause skin diseases and infections.

### Conflict of Interest: None

Ethical Clearance: Not required in an in-vitro studys

### Source of Funding: Self

Acknowledgement: Author Acknowledge the great help received from the laboratory microbiology and laboratory technology of animal product, Faculty of Animal Science, Andalas University and special thanks for Prof. Endang Purwati PhD.

### References

- Hafez, M. M., Maghrabi, I. A., and Zaki, N. M. Toward an Alternative Therapeutic Approach for Skin Infections: Antagonistic Activity of Lactobacilli Against Antibiotic-Resistant aureus Staphylococcus and Pseudomonas aeruginosa. 2013.
- Nwanodi, O. Skin Protective Nutraceuticals: The Current Evidence in Brief, 2, 2018.
- 3. Pieters, N., Brunt, J., Austin, B., and Lyndon, A. R. Efficacy of in-feed probiotics against Aeromonas bestiarum and Ichthyophthirius multifiliis skin infections in rainbow trout (Oncorhynchus mykiss, Walbaum ), 2008; Vol 105: pg 723-732.
- 4. Purwati, E. S. N. Aritonang, S. Melia, I. Juliyarsi and H. Purwanto. Manfaaat Probiotik Bakteri Asam Laktat Dadiah Menunjang Kesehatan Masyarakat. Lembaga Leterasi Dayak (LLD). Padang. 2016.
- 5. Awasthi, S., Rothe, M. J., & Eichenfield, L. F. Atopic dermatitis: Kids are not just little people. Clinics in Dermatology, 2020; Vol 33(6): 594-604.
- Nakatsuji, T., & Gallo, R. L. The role of the skin microbiome in atopic dermatitis. Annals of Allergy, Asthma and Immunology, 2020; Vol122(3): 263-
- 7. Weber, S., Mawdsley, E., & Kaye, D. Antibac terial Agents in the Elderly. 2020.
- Sikorska, H., & Smoragiewicz, W. International

- Journal of Antimicrobial Agents Role of probiotics in the prevention and treatment of meticillinresistant Staphylococcus aureus infections. International Journal of Antimicrobial Agents, 2020; 42(6): 475-481.
- Creech, C. B., Al-zubeidi, D. N., & Fritz, S. A. Prevention of Recurrent Staphylococcal Skin Infections, 2015; Vol 29: 429-464.
- 10. Purwati, E., S. Syukur, dan Z. Hidayat. Lactobacillus sp. Isolasi dari Bivicophitomega sebagai Probiotik. Di dalam Proceeding Lembaga Ilmu Pengetahuan Indonesia, Jakarta. 2005.
- 11. Melia, S., Purwati, Yuherman, Jaswandi, Aritonang, S. N., Purwanto, H. Supernatan Bebas Sel Lactobacillus fermentum L23 yang berpotensi sebagai Antimikroba. No. Paten IDS000002471. Dirjen Kekayaan Intelektual, Kementerian Hukumn dan HAM. 2019.
- 12. Syukur, S., Yolanda, S., and Fachrial, E. Isolation, antimicrobial activity and bioremediation of heavy metal Cadmium (Cd) by using lactic acid bacteria from Dadih Origin Lareh Sago Halaban, Journal of Chemical and Pharmaceutical Research, 2015; 7(9): 235-241.
- 13. Harun, H., Wirasti, Y., Purwanto, B., and Purwati, E. Characterization of lactic acid bacteria and determination of antimicrobial activity in dadih from Air Dingin Alahan Panjang district, Solok regency-West Sumatera. Systematic Reviews in Pharmacy, 2020; Vol 11(3): 583-586.
- 14. Melia, S., Purwati, E., Yuherman, Jaswandi, Aritonang, S. N., and Silaen, M. Characterization of the Antimicrobial Activity of Lactic Acid Bacteria Isolated from Buffalo Milk in West Sumatera (Indonesia) Against Listeria monocytogenes. Pakistan Journal of Nutrition, 2017; Vol 16(8): 645-650.
- 15. FAO/WHO. Joint Expert Consultation on Eavaluation of Gealth and Nutrional Properties of Probiotics in Food Including Powder Mlk With Live Lactic Acid Bacteria. 2001.
- 16. Ray, B dan A. Bhunia. Fundamental Food Microbiology. 3rdEd. Florida. CRC Press. London. New York. 2004
- 17. Juliyarsi, I., Hartini, P, Y., Djamaan, A, Purwanto, H and Purwati, E. Characterization of Lactic Acid Bacteria and Determination of Antimicrobial Activity in Tempoyak from Padang Pariaman

- District, West Sumatra, Indonesia. Pakistan Journal of Nutrition, 2018; Vol 17(10): 506-511.
- 18. Shokryazdan, P., Sieo, C. C., Kalavathy, R., Liang, J. B., Alitheen, N. B., Jahromi, M. F., and Ho, Y. W. Probiotic Potential of Lactobacillus Strains with Antimicrobial Activity against Some Human Pathogenic Strains. BioMed Research International. Vol 2014: 16 pg.
- 19. Halder, D., Mandal, M., Chatterjee, Shiv, S., Pal, Nishith, K., and Mandal, S. Indigenous Probiotic Lactobacillus Isolates Presenting Antibiotic like

- Activity against Human. Biomedicines Journal, 2017; Vol 5(31).
- 20. Mandal, S., and Halder, D. Exploring Anti-Klebsiella pneumoniae Activity of Probiotic Lactobacilli of Curd Origin, 2018; Vol 1(4): 49-53.
- 21. Prabhurajeshwar, C., & Chandrakanth, R. K. ScienceDirect Probiotic potential of Lactobacilli with antagonistic activity against pathogenic strains: An in vitro validation for the production of inhibitory substances. Biomedical Journal, 2017; Vol 40(5): 270-283.

# Antimicrobial Activity of Lactic Acid Bacteria Found in Dadiah on Disease-Causing Skin Infections

### **ORIGINALITY REPORT**

15% SIMILARITY INDEX

12%
INTERNET SOURCES

6%
PUBLICATIONS

% STUDENT PAPERS

**PRIMARY SOURCES** 



repository.unair.ac.id

Internet Source

12%

2

Rinita AMELIA, Koshy PHILIP, Yudha Endra PRATAMA, Endang PURWATI.

3%

"Characterization and probiotic potential of lactic acid bacteria isolated from dadiah sampled in West Sumatra", Food Science and Technology, 2020

Publication

Exclude quotes

On

Exclude matches

< 3%

Exclude bibliography O